
# Psychology of Perception Psychology 4165, Summer 2008 Laboratory 3 Hue-Naming Functions



Lewis O. Harvey, Jr.–Instructor Rosi Kaiser–Assistant Vyga Kaufmann –Assistant MUEN D156, 09:15–10:50 M–F

Lab 3: Hue Naming

Page intentionally blank

Lewis O. Harvey, Jr.–Instructor Rosi Kaiser–Assistant Vyga Kaufmann –Assistant MUEN D156, 09:15–10:50 M–F

# Lab 3: Hue Naming

### Introduction

Color is a psychological experience composed of at least three psychological dimensions: **hue**, **saturation**, and **brightness**. Hue is the experience that we describe with color name labels such as red or blue. Saturation describes the intensity of the hue experience, ranging from hueless to deeply saturated. Pink, for example, is a desaturated red. Brightness is the dimension of experience that permits us to order colors from dark to light. Yellow usually appears brighter than navy blue. In this experiment you will explore the color dimension of hue and gain some insight into the perceptual mechanisms that create it.

Physical light is composed of discrete packets of energy called quanta. A quantum of light has only a single physical property that can be described in one of three ways: By the energy in the quantum; by the frequency of the quantum; or by the wavelength of the quantum. Quanta having wavelengths in the range of approximately 400 to 700 nm (1 nanometer =  $10^{-9}$  meter) are called photons because, under the right circumstances, they can lead to visual experiences. Light has no color; color is an experience created by the visual system in response to stimulation by light.

There are hundreds of different color names in English and in most languages. In spite of the plethora of color words, people with normal color vision can describe almost all colors as being composed of various percentages of red, yellow, green, and blue. For example, one might describe orange as being 60% red and 40% yellow or lime as 70% green and 30% yellow and so on. In this experiment you are going to view monochromatic lights (light composed of a single wavelength) and estimate the percentage of red, yellow, green, and blue making up the color experience you have of each light.

### Methods

**Procedure**: Form groups of 2 or 3. Each will view monochromatic light projected on a white screen by a Bausch and Lomb monochrometer. Thirty-one separate wavelengths ranging from 400 nm to 700 nm in 10 nm steps will be viewed. These wavelengths should be presented in a random order. Record the size of the colored spot and your viewing distance from it. Viewing distance should remain constant throughout the experiment.

Color Naming Procedure: Present each of the 31 wavelengths one at a time. On each trial, write down on your data sheet (see Appendix I) what percentage of the color experience evoked by the wavelength is red. Write down 0 percent if there is no red in your experience. This task may seem very strange at first. You might want to practice a bit before you start in earnest. After all 31 wavelengths have been judged for their redness, repeat the process for yellow, then for green and finally for

Lewis O. Harvey, Jr.–Instructor Rosi Kaiser–Assistant Vyga Kaufmann –Assistant MUEN D156, 09:15–10:50 M–F

### Lab 3: Hue Naming

blue. Be sure to judge all 31 wavelengths for a given color before moving on to the next color. Use a different random order of the wavelengths for each color judgment.

**Pure Color Procedure**: After you have collected the color naming data above, use the method of adjustment to determine the wavelength that gives you the psychological pure colors of blue, green, and yellow. If you have a dichromatic color deficiency, find the wavelength that appears hueless or neutral white or gray. Record these wavelengths on your data sheet in Appendix II. Make at least 10 trials per color (i.e., blue, green, and yellow). Compute the mean wavelength for each color and enter them on the group data sheet. These means can be computed in R (see below).

Individual Data Analyses: Download the Lab3\_Tools file. Double click on the Excel file "Hue\_Name\_Template.txt" and transfer your data from the table in Appendix 1 to the appropriate column. Note that the last three columns will each have only 10 numbers (the pure color wavelengths from Appendix II), not 31 like the other columns. Save the data in a text file with your own name.

The first step in the data analysis is to compute the opponent responses by subtracting the percent green from the percent red and by subtracting the percent blue from the percent yellow. Also compute the total percentage at each wavelength. If red, yellow, green, and blue are sufficient to describe all color experiences, these totals should be 100 percent.

The second step is to compute the mean wavelength giving pure blue, pure green and pure yellow. These steps may be accomplished by the following R code:

```
df <- cbind(df, red_green = df$red - df$green)
df <- cbind(df, yellow_blue = df$yellow - df$blue)
df <- cbind(df, total = df$red + df$yellow + df$green + df$blue)

# Compute the mean wavelengths of the pure colors
pure.blue <- mean(df$pure_blue, na.rm = TRUE)
pure.green <- mean(df$pure_green, na.rm = TRUE)
pure.yellow <- mean(df$pure_yellow, na.rm = TRUE)</pre>
```

Make two sets of graphs presenting your data: Wavelength should be plotted on the abscissa and percent on the ordinate. Plot your red and green data on one graph and your yellow and blue data on a second graph. The R code for producing these graphs is given in the appendix. These two graphs should look like the ones on the next page (Figure 1).


Now plot the two chromatic processes represented by the difference between red and green and between yellow and blue. Use an 8<sup>th</sup> order polynomial to fit a smooth function through your data. Plot the data and the smooth function. Add to

Lewis O. Harvey, Jr.–Instructor Rosi Kaiser–Assistant Vyga Kaufmann –Assistant MUEN D156, 09:15–10:50 M–F

# Lab 3: Hue Naming

this graph vertical lines to mark the position of the wavelength corresponding to pure blue, green, and yellow. Your graphs should resemble those on the next page (Figure 2). The R code to do the polynomial fit and to plot the data is given in the Appendix.

From your chromatic process curves in Figure 2, determine the wavelengths giving psychologically pure blue (where red and green are zero), green (where blue and yellow are zero), and yellow (where red and green are zero). You may use the R functions defined in the script (predict.rg(nm) and predict.yb(nm)) to try different wavelength values to get close to zero. How do these wavelengths compare with those measured directly? In your discussion relate your findings to the opponent-process theory of color vision.



Compare your data (graph) with those of your group members. How are they alike and how do they differ? If you have someone in your group who has a color deficiency examine his (most likely it would be a male) data carefully and compare them with the members of your group who have normal trichromatic color vision.

Now prepare a third figure plotting the total percentage response against wavelength. Use a linear regression (lm() in R) to compute the best-fitting line. Is the slope of this line zero? Are there wavelengths where having only four color names might not be sufficient to describe your experiences?

Lewis O. Harvey, Jr.–Instructor Rosi Kaiser–Assistant Vyga Kaufmann –Assistant MUEN D156, 09:15–10:50 M–F

# Lab 3: Hue Naming

Group Data Analysis: The data file lab3\_data.txt contains the mean wavelengths corresponding to psychologically pure blue, pure green, and pure yellow for each observer in the class. Also are the neutral point wavelengths that should correspond to the same colors. Examine the data to see if there is much consensus among observers. You could plot the wavelengths for the three colors on a strip chart or make a box and whiskers plot to look at the spread of your data. You could also look at the means and standard deviations for each color. Finally, test the hypothesis that the wavelengths giving pure blue, pure green and pure yellow are the same (the null hypothesis). In addition test the null hypothesis that gender makes no difference on the wavelengths giving pure colors. The series of commands in R that create these graphs and compute the statistics are given at the end of this handout.

# **Laboratory Report**

Your lab report should contain five parts: *Cover Sheet, Introduction, Methods, Results,* and *Discussion*. In the *Introduction* explain why you did the experiment. In the *Methods* section describe what you did. In the *Results* section present your findings, including graphs of your data. In the *Discussion* of your results, here are some important questions to answer. Is there a systematic relationship between wavelength of light and the percentages of red, yellow, green, and blue experience evoked by it? Are there any points in the wavelength spectrum that give rise to a unique hue? A unique hue would occur at a wavelength that gave 100% of one color name and 0% of the other three. Look at the relationship among the four curves. Do pairs of curves seem to have a special relationship with one another? Relate your findings to the trichromatic theory of color vision proposed by Helmholtz and to the opponent-process theory proposed by Hering. Examine the spread of the unique hue wavelength across the students in the class. Is there good agreement or not?

Your lab report should be brief and contain six sections: cover page, introduction, methods, results, discussion, and references. These sections should conform to the American Psychological Association (APA) style as described in Chapter 13 of the Martin book. The results section should contain the graphs plotting your data and the opponent-process transformation. The report is due at lab meeting (20 & 21 June 2008). Late labs will receive a grade of zero. All lab reports must be prepared with a word processor. It is worth 50 points.

Lewis O. Harvey, Jr.–Instructor Rosi Kaiser–Assistant Vyga Kaufmann –Assistant MUEN D156, 09:15–10:50 M–F

# Lab 3: Hue Naming

Appendix I: Percent Color Experience

	Appendix I: Percent Color Experience						
	Wavelength	Red	Yellow	Green	Blue		
1	400 nm						
2	410 nm						
3	420 nm						
4	430 nm						
5	440 nm						
6	450 nm						
7	460 nm						
8	470 nm						
9	480 nm						
10	490 nm						
11	500 nm						
12	510 nm						
13	520 nm						
14	530 nm						
15	540 nm						
16	550 nm						
17	560 nm						
18	570 nm						
19	580 nm						
20	590 nm						
21	600 nm						
22	610 nm						
23	620 nm						
24	630 nm						
25	640 nm						
26	650 nm						
27	660 nm						
28	670 nm						
29	680 nm						
30	690 nm						
31	700 nm						

-7-

Lewis O. Harvey, Jr.–Instructor Rosi Kaiser–Assistant Vyga Kaufmann –Assistant MUEN D156, 09:15–10:50 M–F

# Lab 3: Hue Naming

# Appendix II: Psychologically Pure Colors

	Wavelength for psychologically pure color				
Trial	Blue	Green	Yellow		
1					
2					
3					
4					
5					
6					
7					
8					
9					
10					
Mean					

# Appendix III: Opponent-Process Neutral Points from curve-fitting

	Wavelength of Opponent-Process Neutral Points			
Zero point	Blue	Green	Yellow	
nm				

```
# lab3_individual_analysis.R
# command file for analyzing individual hue naming data
# Lewis O. Harvey, Jr.
# Department of Psychology
# University of Colorado
# 10 June 2008
# read file into data frame
df <- read.delim(file.choose(), header=TRUE)</pre>
# Compute opponent processs and add columns to data frame
df <- cbind(df, red_green = df$red - df$green)</pre>
df <- cbind(df, yellow_blue = df$yellow - df$blue)</pre>
df <- cbind(df, total = df$red + df$yellow + df$green + df$blue)</pre>
# Compute the mean wavelengths of the pure colors
pure.blue <- mean(df$pure_blue, na.rm=TRUE)</pre>
pure.green <- mean(df$pure_green, na.rm=TRUE)</pre>
pure.yellow <- mean(df$pure_yellow, na.rm=TRUE)</pre>
par(ask = TRUE)
par(mfrow=c(2,1)) # two graphs per figure
# -----
# plot the individual color percentages
# -----
# red and green percentages
plot(red \sim wavelength, data = df,
    col = "red", type = "b", pch = 19,
    xlim = c(400, 700),
    ylim = c(0, 100),
    xlab = "Wavelength in nm",
    ylab = "Percentage of Color",
    main = "Fig 1: Red and Green Responses")
points(green ~ wavelength, data = df,
    col = "green", type = "b", pch = 19,
    xlim = c(400, 700),
    ylim = c(-100, 100))
# yellow and blue percentages
plot(yellow ~ wavelength, data = df,
    col = "orange", type = "b", pch = 19,
```

```
xlim = c(400, 700),
    ylim = c(0, 100),
    xlab = "Wavelength in nm",
    ylab = "Percentage of Color",
    main = "Fig 1: Yellow and Blue Responses")
points(blue ~ wavelength, data = df,
    col = "blue", type = "b", pch = 19,
    xlim = c(400, 700),
    ylim = c(-100, 100))
# -----
# plot the opponent processes
# -----
# plot the red-green process
df.red <- subset(df, red_green > 0)
df.green <- subset(df, red_green <= 0)</pre>
plot(red_green ~ wavelength, data = df.red,
    type = "p", col = "red", pch = 19,
    xlim = c(400, 700),
    ylim = c(-100, 100),
    xlab = "Wavelength in nm",
    ylab = "Red - Green Response",
    main = "Fig 2: Red - Green Chromatic Mechanism")
points(red_green ~ wavelength, data = df.green,
    type = "p", col = "green", pch = 19)
abline(h = 0)
# fit a polynomial to the opponent process data and
# plot the best-fitting line
rg.poly <- lm(red_green ~ poly(wavelength, degree = 8), data = df)
x <- 400:700
lines(x, predict(rg.poly, data.frame(wavelength = x)))
abline(v = pure.blue, col = "blue")
abline(v = pure.yellow, col = "orange")
# plot the yellow-blue opponent process
df.yellow <- subset(df, yellow_blue > 0)
df.blue <- subset(df, yellow_blue <= 0)</pre>
plot(yellow_blue ~ wavelength, data = df.yellow,
    type = "p", col = "orange", pch = 19,
    xlim = c(400, 700),
    ylim = c(-100, 100),
```

```
xlab = "Wavelength in nm",
    ylab = "Yellow - Blue Response",
    main = "Fig 2: Yellow - Blue Chromatic Mechanism")
points(yellow_blue ~ wavelength, data = df.blue,
    type = "p", col = "blue", pch = 19)
abline(h = 0)
# fit a polynomial to the opponent process data and
# plot the best-fitting line
yb.poly <- lm(yellow_blue ~ poly(wavelength, degree = 8), data = df)
x <- 400:700
lines(x, predict(yb.poly, data.frame(wavelength = x)))
abline(v = pure.green, col = "green")
# define the functions for finding the roots
# of the opponent process equations
predict.rg <- function(x){predict(rg.poly, data.frame(wavelength = x))}</pre>
predict.yb \leftarrow function(x)\{predict(yb.poly, data.frame(wavelength = x))\}
# plot total percentage as a function of wavelength
plot(total \sim wavelength, data = df, pch = 19,
    ylim = c(0, 100),
    xlab = "Wavelength in nm",
    ylab = "Total Percentage",
    main = "Sum of Red, Yellow, Green, and Blue Percentages")
# compute a linear regression on total
tot.lm <- lm(total \sim wavelength, data = df)
lines(tot.lm$fitted ~ df$wavelength)
# plot the residual from the total ~ wavelength regression
plot(tot.lm\residual \sim df\wavelength, pch = 19,
    ylim = c(-50, 50),
    xlab = "Wavelength in nm",
    ylab = "Residuals")
abline(h=0)
par(mfrow=c(1,1))
par(ask = FALSE)
```

```
# lab3_group_analysis.R
# Analysis of Group Hue Namina Data
# 18 June 2008
# load the nlme library to use lme()
library(nlme)
# ask before plotting
par(ask=TRUE)
# plot two graphs in one panel
par(mfcol = c(1,2))
# read in group data in wide format from Excel text file
fn <- file("lab3_group_data_wide.txt")</pre>
df.wide <- read.delim(fn, header=TRUE)</pre>
# *********************
# reshape the file into long format
# Note: reshape() is a bit of a pain to use but what it does
# is take data that are in a so-called wide format
# (one row per subject, separate columns per dependent variable)
# and put it into the long formate needed by many R analysis routines
# (multiple rows per subject, single column per dependent variable)
df.long <- reshape(df.wide,</pre>
    varying = list(c("zeroB", "zeroG", "zeroY"), c("pureB", "pureG",
"pureY")),
    idvar = "subj",
    times = c("blue", "green", "yellow"),
    v.names = c("zero_wavelength", "pure_wavelength"),
    timevar = "color".
    direction = "long")
rownames(df.long) <- 1:dim(df.long)[1]</pre>
df.long$color <- factor(df.long$color)</pre>
# use shorter name for the data frame
df <- df.long</pre>
# Make a strip chart of the zero-crossing data
stripchart(zero_wavelength ~ color, data = df, method = "jitter", jitter =
0.03,
    ylab = "zero-crossing wavelength (nm)",
    xlab = "Color", ylim = c(400, 650), vertical = TRUE)
```

```
# Make a box plot of the zero-crossing data:
boxplot(zero_wavelength \sim color, data = df, ylim = c(400, 650),
    ylab = "zero-crossing wavelength (nm)",
    xlab = "Color", ylim = c(400, 650))
# Make a strip chart of the pure-color data
stripchart(pure_wavelength ~ color, data = df, method = "jitter", jitter =
0.03,
    vlab = "pure color wavelength (nm)",
    xlab = "Color", ylim = c(400, 650), vertical = TRUE)
# Make a box plot of the pure-color data:
boxplot(pure_wavelength \sim color, data = df, ylim = c(400, 650),
    ylab = "pure color wavelength (nm)",
    xlab = "Color", ylim = c(400, 650))
par(mfcol = c(1,1))
# plot the wavelengths for the pure colors from settings
# against the wavelengths derived from zero crossings
par(pty = "s") # make graph square
plot(pure_wavelength ~ zero_wavelength, data = df, type="n",
    xlim = c(400, 650),
    ylim = c(400, 650),
    xlab = "zero-crossing wavelengths",
    ylab = "pure-color wavelength",
    main = "Hue-Naming Laboratory Group Data")
abline(0,1)
points(pure_wavelength ~ zero_wavelength,
data=subset(df,df$color=="blue"), col="blue", pch=19)
points(pure_wavelength ~ zero_wavelength,
data=subset(df,df$color=="green"), col="green", pch=19)
points(pure_wavelength ~ zero_wavelength,
data=subset(df,df$color=="yellow"), col="orange", pch=19)
par(pty = "m")
# Testing the hypothesis that color is independent of wavelength and
aender
# Compute the mixed-effects (within and between factors ANOVA
a \leftarrow lme(pure\_wavelength \sim color * gender, random = \sim 1 | subj, data = df)
# print a summary of the analysis of variance
```

Lewis O. Harvey, Jr.–Instructor Rosi Kaiser–Assistant Vyga Kaufmann –Assistant MUEN D156, 09:15–10:50 M–F

```
print(anova(a))
# Print a table of means for the three colors
print("color means
")
print(tapply(df$pure_wavelength, df$color, mean, na.rm=TRUE))
print(tapply(df$zero_wavelength, df$color, mean, na.rm=TRUE))
# Print a table of means for the genders
print("gender means")
print(tapply(df$pure_wavelength, df$gender, mean, na.rm=TRUE))
print(tapply(df$zero_wavelength, df$gender, mean, na.rm=TRUE))
# Print a table of means for gender and colors
print("gender and color means")
print(tapply(df$pure_wavelength, INDEX=list(df$gender, df$color), mean,
na.rm=TRUE))
print(tapply(df$zero_wavelength, INDEX=list(df$gender, df$color), mean,
na.rm=TRUE))
# Print correlation between the two ways of measuring hue
print("correlation between pure color wavelengths and zero-crossing
wavelengths
")
print(cor(df$pure_wavelength, df$zero_wavelength))
par(ask=FALSE)
```